ADVIL DUAL ACT 25 BX PKG_051723 (Print Ready)

- Drug Facts: 10pt
- Drug Facts (continued): 9pt
- Headings: 8pt
- Subheadings: 6pt
- Bulleted text: 6pt
- Main text: 6pt

Non Varnish Area
No Print

Print Free Zone

Print Free Zone

Print Free Zone

Advil
DUAL ACTION
WITH Acetaminophen

Acetaminophen 250 mg and
Ibuprofen (NSAID) 125 mg Tablets
Pain Reliever

50 CAPLETS
25 packets of 2 caplets each
*Capsule-Shaped Tablets

Advil II

Do not use with other medicines containing ACETAMINOPHEN; can cause liver damage.

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredients (in each caplet)	Purposes
Acetaminophen 250 mg	Pain reliever
Ibuprofen 125 mg (NSAID*)	Pain reliever

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
  - headache ■ toothache ■ backache ■ menstrual cramps ■ muscular aches ■ minor pain of arthritis

Warnings

Acetaminophen liver damage warning:

This product contains acetaminophen. Severe liver damage may occur if you take:
■ with other drugs containing acetaminophen ■ more than 6 caplets in 24 hours, which is the maximum daily amount for this product ■ 3 or more alcoholic drinks every day while using this product

Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.

NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ if you have ever had an allergic reaction to acetaminophen or any other pain reliever ■ right before or after heart surgery

Ask a doctor before use if

■ you have liver disease ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are

■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

Drug Facts *(continued)*

When using this product

■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke:
■ chest pain ■ trouble breathing ■ weakness in one part or side of body
■ slurred speech ■ leg swelling
■ pain gets worse or lasts more than 10 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ **do not take more than directed**

adults and children 12 years and over	**take 2 caplets every 8 hours** while symptoms persist
children under 12 years	ask a doctor

■ do not take more than 6 caplets in 24 hours, unless directed by a doctor

Other information

■ read all warnings and directions before use. Keep carton. ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, ferric oxides, glyceryl dibehenate, hypromellose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, titanium dioxide

Questions or comments?

call weekdays 9 AM to 5 PM at **1-800-88-ADVIL**

Do Not Use if the sealed child resistant pouch is broken or torn.

For most recent product information, visit www.Advil.com. For US Patent or Application status see www.productpats.com. ©2022 GSK group of companies or its licensor. Trademarks are owned by or licensed to the GSK group of companies.

This Product is repackaged by JC World Bell Wholesale Co, Inc; 376-380 Railroad Ave, Hackensack NJ 07601; 201-488-1096 (Mon-Fri 9AM - 6PM EST) with the permission of GSK Consumer Healthcare.

0 73194 6 014058

Glue Area
Non Varnish
Area
Print Free
Zone

Print Free Zone

Print Free Zone

Advil
DUAL ACTION
WITH Acetaminophen

Acetaminophen 250 mg and
Ibuprofen (NSAID) 125 mg Tablets
Pain Reliever

50 CAPLETS
25 packets of 2 caplets each
*Capsule-Shaped Tablets

Advil II

TO OPEN, PUSH IN TAB AND PULL OUT

Do Not Use if the sealed child resistant pouch is broken or torn.

See Full Drug Facts information on the back of this dispenser.

50 CAPLETS (25 PACKETS OF 2 CAPLETS EACH)
*Capsule-Shaped Tablets

Print Free Zone

Print Free Zone

CONTAINS 2 MEDICINES

Acetaminophen + Ibuprofen

Advil
DUAL ACTION
WITH Acetaminophen

Acetaminophen 250 mg and
Ibuprofen (NSAID) 125 mg Tablets
Pain Reliever

Advil II

50 CAPLETS
25 packets of 2 caplets each
*Capsule-Shaped Tablets

Glue Area
Non Varnish
Area
Print Free
Zone

ADVIL ALLERGY 25 BX PKG_052223

- Drug Facts: 9pt
- Drug Facts (continued): 8pt
- Headings: 8pt
- Subheadings: 6pt
- Bulleted text: 6pt
- Main text: 6pt

Print Free Zone

Print Free Zone

Print Free Zone

Print Free Zone

Advil

ALLERGY & CONGESTION RELIEF

Chlorpheniramine Maleate 4 mg Antihistamine
Ibuprofen 200 mg Pain Reliever/Fever Reducer (NSAID)
Phenylephrine HCl 10 mg Nasal Decongestant

1 PILL DOSAGE

Relief of: Sinus Pressure, Nasal Swelling & Congestion, Sneezing, Runny Nose and Headache

Advil Adult

25 POUCHES OF 1 COATED TABLET EACH

Non Varnish Area
No Print

Drug Facts

Active ingredients (in each tablet) — **Purposes**

Chlorpheniramine maleate 4 mg ..... Antihistamine
Ibuprofen 200 mg (NSAID)* ..... Pain reliever/fever reducer
Phenylephrine HCl 10 mg ..... Nasal decongestant

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:
  - runny nose
  - itchy, watery eyes
  - itching of the nose or throat
  - sneezing
  - nasal congestion
  - sinus pressure
  - headache
  - minor body aches and pains
  - fever
- reduces swelling of the nasal passages
- temporarily restores freer breathing through the nose

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning:
NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
- in children under 12 years of age because this product contains too much medication for children under this age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if
- you have a breathing problem such as emphysema or chronic bronchitis
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, glaucoma, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are
- under a doctor's care for any serious condition
- taking sedatives or tranquilizers
- taking any other product that contains phenylephrine, chlorpheniramine or any other nasal decongestant or antihistamine
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product
- take with food or milk if stomach upset occurs
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- may cause excitability especially in children

Drug Facts (continued)

Stop use and ask a doctor if
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- symptoms continue or get worse
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- **do not take more than directed**
- adults and children 12 years of age and over:
  - take 1 tablet every 4 hours while symptoms persist
  - do not take more than 6 tablets in any 24-hour period, unless directed by a doctor
- children under 12 years of age: do not use because this product contains too much medication for children under this age

Other information
- read all warnings and directions before use.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients acesulfame potassium, artificial flavors, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, glyceryl dibehenate, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, silicon dioxide, sucralose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum

Questions or comments?
Call weekdays from **9 AM to 5 PM EST** toll free at **1-800-88-ADVIL**

Do Not Use if the sealed child resistant pouch is broken or torn

For most recent product information, visit www.advil.com. Trademarks owned or licensed by GSK. ©2021 GSK or licensor

This product is repackaged and distributed by JC World Bell Wholesale Co. Inc, 376-380 Railroad Ave, Hackensack NJ 07601, 201-488-1066 (Mon-Fri 9AM - 6PM EST) with the permission of GSK Consumer Healthcare.

0 731946 014003

Glue Area
Non Varnish
Area
Print Free
Zone

Print Free Zone

Print Free Zone

Advil

ALLERGY & CONGESTION RELIEF

TO OPEN, PUSH IN TAB AND PULL OUT

Do Not Use if the sealed child resistant pouch is broken or torn. See Full Drug Facts information on the back of this dispenser.

Chlorpheniramine Maleate 4 mg Antihistamine
Ibuprofen 200 mg Pain Reliever/Fever Reducer (NSAID)
Phenylephrine HCl 10 mg Nasal Decongestant

1 PILL DOSAGE

Relief of: Sinus Pressure, Nasal Swelling & Congestion, Sneezing, Runny Nose and Headache

Advil Adult

25 POUCHES OF 1 COATED TABLET EACH

See Full Drug Facts information on the back of this dispenser.

Print Free Zone

Print Free Zone

Advil

ALLERGY & CONGESTION RELIEF

Chlorpheniramine Maleate 4 mg Antihistamine
Ibuprofen 200 mg
Pain Reliever/Fever Reducer (NSAID)
Phenylephrine HCl 10 mg
Nasal Decongestant

1 PILL DOSAGE

Relief of: Sinus Pressure, Nasal Swelling & Congestion, Sneezing, Runny Nose and Headache

Advil Adult

25 POUCHES OF 1 COATED TABLET EACH

Glue Area
Non Varnish
Area
Print Free
Zone

ALEVE 25's Box PKG_081524 (Print-Ready 082924)

ALKA SELTZER '72'S BOX (DSP BOX)'_032222 PRINT READY

Print Free Zone

Print Free Zone

Print Free Zone

Print Free Zone

Print Free Zone

Alka-Seltzer®

Buffered aspirin (NSAID) / Pain reliever / fever reducer

Fast Relief of:
Headache
Backache
Body Aches
Muscle Pain

BAYER

ORIGINAL FLAVOR

72 EFFERVESCENT TABLETS

Non Varnish Area
No Print

Drug Facts

Active Ingredient (in each tablet) — **Purpose**

Buffered aspirin 325 mg (NSAID)* ..... Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - muscle pain
  - backache
  - toothache
  - menstrual pain
  - colds
  - minor pain of arthritis
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warnings: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- diabetes
- gout
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- fully dissolve 2 tablets in 4 ounces of water before taking

adults and children 12 years and over	2 tablets every 4 hours, or as directed by a doctor	do not exceed 8 tablets in 24 hours
adults 60 years and over	2 tablets every 4 hours, or as directed by a doctor	do not exceed 4 tablets in 24 hours
children under 12 years	consult a doctor	

Other information

- each tablet contains: sodium 568 mg
- store at room temperature. Avoid excessive heat.
- Alka-Seltzer Original Flavor in water contains principally the analgesic sodium acetylsalicylate

Inactive Ingredients anhydrous citric acid, sodium bicarbonate

Questions or comments? 1-800-968-0369 (Mon – Fri 9AM – 5PM EST)

Dist. by: Bayer HealthCare LLC Whippany, NJ 07981

Bayer, the Bayer Cross and Alka-Seltzer are registered trademarks of Bayer.

This product is repackaged by JC world Bell Wholesale Co, Inc; 376-380 Railroad Ave, Hackensack NJ 07601; 201-488-1096 (Mon-Fri 9AM-6PM EST), from a product manufactured by Bayer.

MADE IN MEXICO

LOT: EXP:

8 52682 50847 9

Alka-Seltzer®

ORIGINAL FLAVOR

Fast Relief of:
Headache
Backache
Body Aches
Muscle Pain

BAYER

72 EFFERVESCENT TABLETS

Non Varnish Area Print Free Zone

Alka-Seltzer®

Buffered aspirin (NSAID) / Pain reliever / fever reducer

Fast Relief of:
Headache
Backache
Body Aches
Muscle Pain

BAYER

ORIGINAL FLAVOR

72 EFFERVESCENT TABLETS

TO OPEN
PUSH IN TAB AND PULL OUT

Do not use if pouch is torn or open

72 EFFERVESCENT TABLETS

Alka-Seltzer®

ORIGINAL FLAVOR

- Fully dissolve two (2) tablets in 4 oz of water before taking.
- You do not have to drink any of the residue that may be on the bottom of the glass. The medicine of Alka-Seltzer® is in the water.

Take Alka-Seltzer® at any time- morning, noon, or night- when you need relief from headaches, backaches, body aches or muscle pain.

Non Varnish Area Print Free Zone

Print Free Zone

Print Free Zone

Print Free Zone

Print Free Zone

BAYER 50'S PKG_041022

Claritin 25 BX PKG_040523 (Print-Ready)

EMERGENC 25 BX PKG_071523 (2) print ready

- Drug Facts: 20pt
- Headings: 7pt
- Special Characters: 4.5pt
- Main text: 6pt
- Numbers: 6pt

- Antioxidants...: 14pt
- 1,000: 40pt
- mg: 25pt
- Vitamin C: 36pt
- Daily Immune...: 17pt

Non Varnish Area
No Print

Print Free Zone

Our boxes are printed with vegetable-based inks.

This box is recyclable.

Recyclable

Emergen-C

1,000 mg Vitamin C

Daily Immune Support*

Antioxidants,* B Vitamins & Electrolytes

Directions: Ages 14 and up, one (1) packet daily. Empty contents into a glass, add 4–6 oz of water, stir. For lighter flavor, mix with more water. Do not exceed suggested use. If you are pregnant, nursing, or taking medication, consult your doctor before use.

Supplement Facts

Serving Size 1 Packet (9.1 g)

Amount Per Serving		% Daily Value
Calories	35	
Total Carbohydrate	8 g	3%†
Total Sugars	6 g	*
Includes 6 g Added Sugars		12%†
Vitamin C (as ascorbic acid, zinc ascorbate)	1,000 mg	1,111%
Thiamin (as thiamine hydrochloride) (Vit. B_1)	0.36 mg	30%
Riboflavin (as riboflavin-5'-phosphate) (Vit. B_2)	0.39 mg	30%
Niacin (Vit. B_3)	4 mg	25%
Vitamin B_6 (as pyridoxine hydrochloride)	10 mg	588%
Folate (Vit. B_9)	167 mcg DFE (100 mcg Folic Acid)	42%
Vitamin B_{12} (as cyanocobalamin)	25 mcg	1,042%
Pantothenic Acid (as calcium pantothenate) (Vit. B_5)	2.5 mg	50%
Calcium (as calcium carbonate, monobasic calcium phosphate, tribasic calcium phosphate, calcium pantothenate)‡	50 mg	4%
Phosphorus (as monobasic potassium phosphate, monobasic calcium phosphate, monobasic sodium phosphate, tribasic calcium phosphate)‡	38 mg	3%
Magnesium (as magnesium hydroxide, magnesium carbonate)‡	53 mg	13%
Zinc (as zinc ascorbate)	2 mg	18%
Manganese (as manganese gluconate)	0.5 mg	22%
Chromium (as chromium picolinate)	10 mcg	29%
Sodium (as sodium bicarbonate, monobasic sodium phosphate)†	65 mg	3%
Potassium (as potassium bicarbonate, potassium carbonate, monobasic potassium phosphate)‡	200 mg	4%

†Percent Daily Values are based on a 2,000 calorie diet.
* Daily Value not established. ‡Electrolytes

Other Ingredients: Fructose, Maltodextrin, Citric Acid, Malic Acid. **Contains <2% of:** Acacia, Beta-Carotene (color), Glycine, L-Aspartic Acid, Natural Flavors, Orange Juice Concentrate (color and flavor), Orange Oil, Silicon Dioxide, Tartaric Acid, Tocopherols (to preserve freshness).

Distributed by: AlacerCorp,
Warren, NJ 07059
Questions? Comments? Call 1-800-245-1040
For most recent product information, visit www.emergenc.com
Trademarks are owned by or licensed to the Haleon group of companies.
©2022 Haleon group of companies or licensors.
This Product is repackaged by JC World Bell Wholesale Co, Inc; 376-380 Railroad Ave, Hackensack NJ 07601; 201-488-1096 (Mon-Fri 9AM - 6PM EST)
Caffeine-Free • Gluten-Free • Vegetarian • Natural Sweeteners & Flavors
Keep out of reach of children.

0 731946 013792

*Based on the USDA.gov nutrient database for a large, raw orange

Print Free Zone

Glue Area
Non Varnish Area
Print Free Zone

Immune Support

1,000 mg of vitamin C plus other antioxidants zinc and manganese support your immune system.*

Energy

7 B vitamins including B_1, B_2, B_3, B_5, B_6, B_9 and B_{12} enhance energy naturally* without caffeine.

‡Electrolytes

Great for post-workout, replace key electrolytes lost through perspiration.*

feel the Good

Store at room temperature. Protect from moisture.
Carton contains individual packets.
Do not use if packet seal is torn or broken.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

Print Free Zone

Glue Area
Non Varnish Area
Print Free Zone

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

Emergen-C

1,000 mg Vitamin C

Daily Immune Support*

Antioxidants,* B Vitamins & Electrolytes

Super Orange

Flavored Fizzy Drink Mix with other Natural Flavors

Caffeine-Free, Gluten-Free, Vegetarian & Natural Sweeteners

Natural fruit flavors

As much Vitamin C as 10 oranges!*

DIETARY SUPPLEMENT *Vitamin C, Zinc & Manganese

25 - 0.32 OZ (9.1 g) PACKETS / NET WT. 8 OZ (227.5 g)

25 PACKETS

Print Free Zone

Print Free Zone

Naturally, It's Good for You!

Bursting with all sorts of sunshine-y citrus deliciousness, it's no wonder we call it "Super." With each sip, you can feel the essential nutrients flow through your body in a wave of Emergen-C rejuvenation.

If feeling good is your thing, you found the right box.

Get your free samples at emergenc.com!§

§ While supplies last. See emergenc.com for limitations.

Less Water = More Flavor

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

Glue Area
Non Varnish Area
Print Free Zone

EXCEDRIN MIGRAINE 25's Box PKG_081524 (Print-Ready 082924)

TAMPER-EVIDENT BOTTLE
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active Ingredients (in each caplet)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID*)	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen.
Severe liver damage may occur if you take
• more than 2 caplets in 24 hours, which is the maximum daily amount
• with other drugs containing acetaminophen
• 3 or more alcoholic drinks every day while using this product

Stomach bleeding warnings: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
• are age 60 or older • have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month.

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if
• you have never had migraines diagnosed by a health professional
• you have a headache that is different from your usual migraines
• you have the worst headache of your life • you have fever and stiff neck
• you have headaches beginning after or caused by head injury, exertion, coughing or bending • you experienced your first headache after the age of 50 • you have daily headaches • you have a migraine so severe as to require bed rest • you have liver disease

Drug Facts (continued)
• stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have problems or serious side effects from taking pain relievers or fever reducers • you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are
• taking a prescription drug for: • diabetes • gout • arthritis
• under a doctor's care for any serious condition
• taking any other drug • taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if
• an allergic reaction occurs. Seek medical help right away.
• you experience any of the following signs of stomach bleeding:
  • feel faint • vomit blood • have bloody or black stools
  • have stomach pain that does not get better
• your migraine is not relieved or worsens after first dose
• new or unexpected symptoms occur • ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • **do not use more than directed** • adults: take 2 caplets with a glass of water • if symptoms persist or worsen, ask your doctor • do not take more than 2 caplets in 24 hours, unless directed by a doctor • under 18 years of age: ask a doctor

Other Information store at controlled room temperature 20°–25°C (68°–77°F) • close cap tightly after use • read all product information before using. Keep this box for important information.

Inactive Ingredients benzoic acid, carnauba wax, FD&C blue no. 1, hypromellose, light mineral oil, low-substituted hydroxypropyl cellulose, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone, sorbitan monolaurate, stearic acid, titanium dioxide

Questions or comments? 1-800-488-7746

2024 Haleon or licensor. Trademarks are owned by or licensed to the Haleon group of companies.

LOT : EXP :

This Product is repackaged by JC World Bell Wholesale Co, Inc; 376-380 Railroad Ave, Hackensack NJ 07601; 201-488-1995 (Mon-Fri 9AM - 6PM EST), from a product manufactured by Haleon.

0 73146 01492

Hangover Helper (La Rapidita) 25'S BOX PKG_032024-2



Midol 25's PKG_New (Aug 2022)_081622 (Print-ready)

See New Warnings

Motrin® IB

Ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

Convenient Take-Along Packs

Coated Caplets**

**Capsule-Shaped Tablets

Actual Size

Drug Facts

Active ingredient (in each caplet) **Purposes**

Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better

Drug Facts (continued)

- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**

adults and children 12 years and older	take 1 caplet every 4 to 6 hours while symptoms persist if pain or fever does not respond to 1 caplet, 2 caplets may be used do not exceed 6 caplets in 24 hours, unless directed by a doctor
children under 12 years	ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- **do not use if pouch is torn or damaged**

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, iron oxides, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, stearic acid, talc, titanium dioxide

Questions? Call **1-877-895-3665** (toll-free) or **215-273-8755** (collect) or visit www.motrin.com

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2017
www.motrin.com
30038844

8K0B3 MN C2

Product repackaged and distributed with permission of the manufacturer by:
JC World Bell Wholesale Company, Inc. Hackensack, NJ 07601

8 52682 50643 7

See New Warnings

Motrin® IB

Ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

Convenient Take-Along Packs

Coated Caplets**

**Capsule-Shaped Tablets

Actual Size

TO OPEN PUSH IN TAB AND PULL OUT

Convenient Take-Along Packs

25 Pouches of 2 Coated Caplets** Each
Total 50 Coated Caplets**

**Capsule-Shaped Tablets

See New Warnings

Motrin® IB

Ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

Convenient Take-Along Packs

Actual Size of Pouch

Actual Size

Coated Caplets**

**Capsule-Shaped Tablets

Pepcid 25's_Dispenser box PKG_072320_draft 03

THERAFLU NT (LEMON) 20'S BX PKG_032124 (Print-Ready)

Print Free Zone

Print Free Zone

Print Free Zone

Print Free Zone

Non Varnish Area
No Print

READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE. KEEP CARTON FOR REFERENCE. DO NOT DISCARD.

Drug Facts

Active ingredients (in each packet)	Purposes
Acetaminophen 650 mg	Pain reliever/Fever reducer
Diphenhydramine HCl 25 mg	Antihistamine/Cough suppressant

Uses

- temporarily relieves these symptoms due to a cold:
  - minor aches and pains
  - minor sore throat pain
  - headache
  - runny nose
  - sneezing
  - itchy nose or throat
  - itchy, watery eyes due to hay fever
  - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Do not use
- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are
- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts
- pain or cough gets worse or lasts more than 7 days

These could be signs of a serious condition.

Glue Area
Non Varnish
Area
Print Free
Zone

Print Free Zone

Print Free Zone

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- take every 4 hours, while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

- dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- If using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other information

- **each packet contains:** potassium 10 mg, sodium 23 mg
- **phenylketonurics:** contains phenylalanine 13 mg per packet
- store at controlled room temperature 20-25°C (68-77°F). Protect product from heat and moisture.

Inactive ingredients acesulfame potassium, anhydrous citric acid, aspartame, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, maltodextrin, natural and artificial flavors, silicon dioxide, sodium citrate, soy lecithin, sucrose, tribasic calcium phosphate

Questions or comments? 1-855-328-5259

TAMPER EVIDENT INNER UNIT DO NOT USE IF SEALED THERAFLU PACKET IS TORN OR BROKEN

LOT : **EXP :**

1-855-328-5259

©2023 Haleon group of companies or its licensor. Trademarks are owned by or licensed to the Haleon group of companies. Pat.Infowww.productpats.com

0 731946 015994

This Product is repackaged and distributed by: JC World Bell Wholesale Co, Inc; 376-380 Railroad Ave, Hackensack NJ 07601; 201-488-1096 (Mon-Fri 9AM - 6PM EST), with the permission of Haleon group of companies.

Print Free Zone

Print Free Zone

Glue Area
Non Varnish
Area
Print Free
Zone

TUMS 25's Box PKG_100924-2

Tylenol ES Card package_1 pack_111519_draft 3

FRONT

FRONT CARD

61mm (2.40in) x 72mm(2.83in)

BACK

Tylenol Sinus Inner package & Card_2 pack_111419_draft 3

FRONT

FRONT CARD

61mm (2.40in) x 72mm(2.83in)

BACK



Visine_PKG Box_NEW_071620

VISINE®
RED EYE
HYDRATING
COMFORT
LUBRICANT / REDNESS RELIEVER EYE DROPS

WORKS IN 30 SECONDS
Redness + Irritation Relief

6 COUNT
0.28 FL OZ (8 mL) Bottles

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
Skillman, NJ 08558 Made in Belgium ©J&JCI 2020
Product is repackaged and distributed with permission of
Manufacturer by JC WorldBell Wholesale co. Inc.
376-380 Railroad Ave, Hackensack, NJ, 07601
Office # 1-201-488-1096.

VISINE®
RED EYE
HYDRATING
COMFORT
LUBRICANT / REDNESS RELIEVER EYE DROPS

WORKS IN 30 SECONDS
Redness + Irritation Relief

6 COUNT
0.28 FL OZ (8 mL) Bottles

TO OPEN PUSH IN
AND PULL OUT

6 COUNT
0.28 FL OZ (8 mL) Bottles

VISINE®
RED EYE
HYDRATING
COMFORT
LUBRICANT / REDNESS RELIEVER EYE DROPS

WORKS IN 30 SECONDS
Redness + Irritation Relief

6 COUNT
0.28 FL OZ (8 mL) Bottles

VISINE®
RED EYE
HYDRATING
COMFORT
LUBRICANT / REDNESS RELIEVER EYE DROPS

WORKS IN 30 SECONDS
Redness + Irritation Relief

6 COUNT
0.28 FL OZ (8 mL) Bottles

TEAR HERE
TOP NEEDS TO REMOVED
BEFORE PLACEMENT

VISINE®
RED EYE
HYDRATING
COMFORT
LUBRICANT / REDNESS RELIEVER EYE DROPS

WORKS IN 30 SECONDS
Redness + Irritation Relief

6 COUNT
0.28 FL OZ (8 mL) Bottles

Read directions and warnings before use. Keep this card. It contains important information.

Caution: Do not use if VISINE® imprinted neckband on bottle is broken or missing.

Drug Facts

Active Ingredients — **Purpose**
Polyethylene glycol 400 1% ..... Lubricant
Tetrahydrozoline HCl 0.05% ..... Redness reliever

Uses • for the relief of redness of the eye due to minor eye irritations • for the temporary relief of burning, irritation, and discomfort due to dryness of the eye or exposure to wind or sun • for use as a protectant against further irritation or to relieve dryness of the eye

Warnings
For external use only.
Ask a doctor before use if you have narrow angle glaucoma.

When using this product
- pupils may become enlarged temporarily
- overuse may cause more eye redness
- remove contact lenses before using
- replace cap after each use
- do not use if this solution changes color or becomes cloudy
- do not touch tip of container to any surface to avoid contamination

Stop use and ask a doctor if
- you feel eye pain
- redness or irritation of the eye lasts
- changes in vision occur
- condition worsens or lasts more than 72 hours

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions • adults and children 6 years of age and over: put 1 or 2 drops in the affected eye(s) up to 4 times a day • children under 6 years of age: consult a doctor • children under 2 years of age: do not use

Other Information Store at 20° to 25°C (68° to 77°F)

Inactive Ingredients ascorbic acid, boric acid, dextrose, glycerin, glycine, hypromellose, magnesium chloride, polyquaternium-42, potassium chloride, purified water, sodium citrate, sodium lactate, sodium phosphate dibasic

Questions? call toll-free **888-734-7648** or **215-273-8755** (collect)

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.** Skillman, NJ 08558
Made in Belgium ©J&JCI 2020 www.visine.com
Product is repackaged and distributed with permission of Manufacturer by JC WorldBell Wholesale co. Inc.
376-380 Railroad Ave, Hackensack, NJ, 07601 Office # 1-201-488-1096.

LOT:

EXP:

8 52682 50821 9

NyQuil (Vapocool)_Blister Card PKG 1PC (Resize)_072323

INSERT CARD

Insert Card 2.2750 x 2.90

FRONT

Blister Card 2.9130 x 4.75

BACK

Drug Facts (middle side) 2

Zantac 360 1's_Blister Card file_080521

FRONT

FRONT CARD

61mm (2.40in) x 72mm(2.83in)

BACK



ZYRTEC 20'S BX PKG_060324

'ZiQ' Bandage_PKG_12box (77x82x98 mm)_081220

77x82x98mm

'ZiQ' Bandage_PKG_10pcs (3.7x9.3x1.2)_081220

37x93x12mm

CR-DIGI SCALE (0.01g) CR AP-100 BLK (LED LIGHT)_PKG Design 111522 Print-Ready

Foil Effect
Embossed

CR Digital Scale - AD100 (113x34x172mm) RED - 071921

113x34x172mm

'CR' Digital Scale_DIGIX15X2 RA-500 BLK (BEE)_PKG Box_(142x82x28mm)_0603121

RA 142X82X28mm

88 X 40mm

DIGI SCALE (0.1g) CR 1000 BLK* (NEW LOGO)
061322

(Prism) Hologram
Embossed

97X146X27MM

DIGI SCALE_DIGIX19X4 (0.01g) CR AV500 BLK_081221

AV BOX size:122 x 94 x 32mm

Airssassin

bottle design sample 3PKG_053121.ai

sample mockup

Airssassin_aquapurple_bottle 30ml PKG_070821 Print

Airssassin_Spray 30ml AQUA PURPLE Box 'PDQ BOX' Diecut_070821

PDQ Box

Oral-B Glide_Dispenser 1's Card PKG_012121_04

Oral-B Blister Card Size change - same as Visine size

FRONT

BACK

ORAL-B GLIDE_PACKAGE BOX_121220_03

NATIONWIDE VITAMINS_ADULT SEVEN

Inner Box package_030920_draft 7

Length: 3.0625"(3-1/16") / width: 7" / height: 4.25"(4-1/4")

NATIONWIDE VITAMINS_ADULT SEVEN

Card package_030920_draft 7

width: 2.875" / height: 4.1"

ZiQ_Pregnancy Test_PKG Box(132x60mm)_090420

sample_mockup

'Abelia' sanitizer_lable PKG_500ml(16oz)_GLE_052820



Front

Back

Abelia Lable (GEL)
W: 215 mm, H: 115 mm

Front

Back

sample mockup
(BG : white color)

'Abelia' sanitizer_MIST-lable PKG

‘ABELIA’ _WET WIPES_PKG_WIPEX1X1 - WIPES ANTIBACTERIAL 20'S



package mockup (DO NOT PRINT)

‘ABELIA’ _WET WIPES_PKG_WIPEX1X3 - WIPES ANTIBACTERIAL 100'S

package mockup (DO NOT PRINT)

WeEZ_Glass Jar PKG design 'Black'_110521

Top 159*104*79mm

bottom 155*99*77mm



'CR' Butane Gas_90ml_Display 45x117mm _082520



Mockup (Sample Images)

'CR' Butane Gas_300ml_PKG bottle (52 x 195)_072720

Mockup - Front



Mockup - Back

Sample

'CR' Butane Gas_75ml_PKG bottle (35x135mm)_080620-1

Package Color
- Yellow : PANTONE P 10-8 C
- Black : PANTONE P Process Black C
- Grey : PANTONE P 179-3 C
- Blue : PANTONE P 99-8 C
- Red : PANTONE P 53-8 C

Mockup - Front

Mockup - Back

Sample

Cigar Cutter Inner box (15.5cm x 9.2cm x 3.8cm)



◆ Snap lock base, counter display box is delivered to the customer the way that bottom end is glued and folded. It is already setup and ready to use when unfolded.

◆ It is also known as display box.

◆ It enables quick setup and is suitable for both automatic and manual filling.

◆ Snap lock base, counter display box works well for heavier products. The product should be maximum 1 kg. and shouldn' t spill or leak. If the product is dusty or liquid, it should have its own packaging and the product shouldn' t contact the interior of box directly.

Hair Combs
Made in China
Distributed by NY CATS
0 731946 013747
ZIQ HAIR COMBS 72'S POCKET BLK
Hair Combs
ZIQ HAIR COMBS
72'S POCKET BLK

Hair Combs
Made in China
Distributed by NY CATS
0 731946 013747
ZIQ HAIR COMBS 72'S POCKET BLK
Hair Combs
ZIQ HAIR COMBS
72'S POCKET BLK

PRINT READY 072324

ZIQ - DURAG SILKY SATIN BLACK

Product Dimension W*H : 4.88*8.56" / 0.56*7.69"

3 types > Black, white, red, navy

Quantity -- pc/pk (1 each)

LOGO:
FOIL STICKERS





PRINT READY 072324

ZIQ - DURAG TIE DOWN BLACK

LOGO:
FOIL STICKERS

Product Dimension W*H : 4.69*7"
3 types > Black, white, red, navy
Stocking and tiedown have the same dimensions. The only difference is the quantity. (Stocking is 2 pcs/pk, tiedown is 1 pc/pk)



Cigar Cutter Inner box (16cm x 9.2cm x 3.8cm)

JC WORLD BELL_LIGHTER LEASH_DESIGN 050921

SAMPLE MOCKUP DESIGN



SAMPLE DESIGN GUIDELINE

COLOR SCHEME

LockMan _LOCKX1X1_20mm_2PCS_Card_013020_draft 2

FRONT



BACK



2020 Regi Roll Coreless_50PC_2.25" 50FT, 2.25" 80FT, 3.125" 230FT_071323 (print ready)

Regi Roll Coreless
2.25' 50FT BPA FREE
50PC /CS



10pc package

Pantone 198C

Regi Roll Coreless
2.25' 80FT BPA FREE
50PC /CS



10pc package



Pantone 285C

Regi Roll Coreless
3.125" 230FT BPA FREE
50PC /CS



10pc package

Pantone 2593C

YORIENN CHICKEN

KOREAN FRIED CHICKEN & YORI SHOP

TO-GO PACKAGE BOX DESIGN